

1. Open Dot Experiments

2. Kondo effect


3. Few Electron Dots

**4. Double Quantum Dots**

van der Wiel et al., RMP75, 1 (2003)

A. C. Johnson, Ph. D. Thesis (2005)

# Double Quantum Dots




mutual charging energy

$$E_m = \frac{e^2}{C_m} \left( \frac{C_1 C_2}{C_m^2} - 1 \right)^{-1}$$

interdot tunneling  $t$

$$G_m = 4\pi \frac{e^2}{h} \left( \frac{t}{\Delta} \right)^2$$




$t < \Delta$  well localized electrons

individual charging energies

$$E_{c1(2)} = \frac{e^2}{C_{1(2)}} \left( 1 - \frac{C_m^2}{C_1 C_2} \right)^{-1}$$

# Double Quantum Dots: Quadruple Points




$$E_m = \frac{e^2}{C_m} \left( \frac{C_1 C_2}{C_m^2} - 1 \right)^{-1} \rightarrow 0$$

costs zero energy to add a 2nd electron to other dot if one electron is already present


$$E_{C1(2)} = \frac{e^2}{C_{1(2)}} \quad \text{individual charging energies}$$

assume well localized electrons (weak tunneling, but large enough to measure a current)

- quadruple points  
degeneracy of four charge states





# Double Quantum Dots: Triple Points and Honeycombs




$$0 < C_m < C_{1,2} \quad 0 < E_m < E_{C_1, C_2}$$

(1,1) – (0,0) degeneracy lifted

quadruple points split into two triple points




hole like process



electron like process

## Double Quantum Dots: Single Dot Limit

---




$$0 < C_m \sim C_{1,2}$$



$$E_m \sim E_{C_1, C_2}$$

double dot behaves like a  
single dot with two plunger gates

# Double Quantum Dots



# Double Dot Experiment




# Double Dot Hamiltonian

$$\begin{aligned}
 H_{DQD} = & \frac{E_{c1}}{2} N(N-1) - \frac{NE_{c1} + ME_m}{e} (C_{g1}V_{g1} + C_sV_s) + \sum_{i,\sigma} N_{i\sigma} \epsilon_{i\sigma} \\
 & + \frac{E_{c2}}{2} M(M-1) - \frac{ME_{c2} + NE_m}{e} (C_{g2}V_{g2} + C_dV_d) + \sum_{j,\sigma} M_{j\sigma} \epsilon_{j\sigma} \\
 & + E_m NM + \sum_{i,j,\sigma} t_{ij\sigma} (c_{i\sigma}^\dagger c_{j\sigma} + h.c.) + \text{lead tunneling} \quad (3.11)
 \end{aligned}$$

individual charging
electrostatic
quantum confinement

mutual charging
inter-dot tunneling




electrons well localized

$$G_m < e^2/h$$




# Double Dot Capacitances in the Honeycombs



$$\Delta V_{g1} = \frac{|e|}{C_{g1}}$$

$$\Delta V_{g2} = \frac{|e|}{C_{g2}}$$



$$\Delta V_{g1}^m = \frac{|e|C_m}{C_{g1}C_2} = \Delta V_{g1} \frac{C_m}{C_2}$$


$$\Delta V_{g2}^m = \frac{|e|C_m}{C_{g2}C_1} = \Delta V_{g2} \frac{C_m}{C_1}$$

$\Delta V$  : detuning  
controls energy difference  $\Delta$   
between the dot levels  
keeping constant the  
total dot occupation  $N + M$



# Double Dot Transport

triple points:  
sequential tunneling


honey comb lines:  
cotunneling






# Double Dot Experiment



finite bias: nonlinear transport






# Double Dot at finite bias: Excited State Spectroscopy




triple points expands into triangles obeying  $0 \leq \mu_1 \leq \mu_2 \leq eV$

# Double Dot Experiment: Finite Bias



# Interdot Tunneling: Anticrossing




$$\mathbf{H}_0|\phi_1\rangle = E_1|\phi_1\rangle$$

$$\mathbf{H}_0|\phi_2\rangle = E_2|\phi_2\rangle$$

$$\mathbf{T} = \begin{pmatrix} 0 & t_{12} \\ t_{21} & 0 \end{pmatrix}, \quad t_{12} = t_{21}^*, \quad t_{21} = |t_{21}|e^{i\varphi}$$

$$\mathbf{H} = \mathbf{H}_0 + \mathbf{T}$$



$$\mathbf{H}|\psi_B\rangle = E_B|\psi_B\rangle$$

$$\mathbf{H}|\psi_A\rangle = E_A|\psi_A\rangle$$

$$E_B = E_M - \sqrt{\frac{1}{4}(\Delta E)^2 + |t_{12}|^2}$$

$$E_A = E_M + \sqrt{\frac{1}{4}(\Delta E)^2 + |t_{12}|^2}$$